




Using LSST to Probe the Fundamental Nature of Dark Matter

...OR...

Alex Drlica-Wagner
LSST Cross-Correlation Workshop
May 25, 2016








A Jam Session about Multi-Wavelength Multi-Messenger Synergistic Cross-Correlations for Probing the Nature of Dark Matter


Alex Drlica-Wagner
LSST Cross-Correlation Workshop
May 25, 2016



Observable Universe



Galaxy Superclusters



Galaxy Clusters



Galaxies



 $\sim 10^{23} \ M_{\odot}$




 $\sim 10^{15}\, M_{\odot}$

 $\sim 10^{13} \, \mathrm{M} \, \odot$


Dwarf Spheroidal Galaxies




Black Holes as MACHOs




Minimum Microhalo Mass




WIMP Dark Matter




~10 8 M $_{\odot}$ ~10 $^{-6}$ M $_{\odot}$ ~10 $^{-55}$ M $_{\odot}$





"Pillars" of Dark Matter







Direct Detection



Differential Event Rate

(signal in data)

$$\frac{dN}{dE_r}$$


Particle & Nuclear Physics

(cross section & form factor)

$$\frac{\sigma_0}{2\mu^2 m_{\rm DM}} F(q)^2$$



$$\rho_{\rm DM} \int_{v_{\rm min}}^{v_{\rm esc}} \frac{f(v)}{v} dv$$












Local Dark Matter Distribution

(line-of-sight integral)









Direct Detection








Local Dark Matter Density






year




Local Dark Matter Velocity







Mao et al. (2012) [1210.2721]



Direct Detection







Indirect Detection



Gamma-rays

Anti-matter

Particle Flux

(signal in data)

$$\frac{d\Phi_{\gamma}}{dE_{\gamma}}(E_{\gamma},\phi,\theta) =$$




(photons per annihilation)

$$\frac{d\Phi_{\gamma}}{dE_{\gamma}}(E_{\gamma},\phi,\theta) = \frac{1}{4\pi} \frac{\langle \sigma_{ann} v \rangle}{2m_{WIMP}^2} \sum_{f} \frac{dN_{\gamma}^f}{dE_{\gamma}} B_f$$

X









$$\int_{\Delta\Omega(\phi,\theta)} d\Omega' \int_{los} \rho^2(r(l,\phi')) dl(r,\phi')$$



















Galactic Center








Galactic Center








Extragalactic Background



Actual Cross-Correlation!

Fermi-LAT & Galaxy Catalogs

Fermi-LAT x CFHTLenS



Cuoco et al. (2015) [1506.01030]


Shirasaki et al. (2015) [1511.07092]

Shirasaki et al. (2014) [1404.5503]



Dwarf Galaxies








"Missing" Satellites?








"Missing" Satellites?










Astrophysical Probes















Self-Interacting Dark Matter



Non-Interacting Dark Matter



Self-Interacting Dark Matter








Self-Interacting Dark Matter












Strong Lensing and "Missing" Satellites




Mass modeling of galaxy-scale lenses

Flux ratios can be used to measure the contribution of substructure


Strong predictions from LCDM








Lovell et al. (2013) [1308.1399]



Dalal & Kochanek (2002) [55152] Vegetti & Koopmans (2009) [0903.4752] Hezavah et al. (2014) [1403.2720]




Too Big to Fail



WDM and SIDM can produce cores in dwarf galaxies

... so can baryonic feedback




Elbert et al. (2014) [1412.1477]

Vogelsberger et al. (2014) [1405.5216]

Lovell et al. (2013) [1308.1399]

Alex Drlica-Wagner | Fermilab



Walker & Peñarrubia (2011) [1108.2404]




Summary



- We would like to understand the fundamental unit of dark matter.
- Our knowledge is currently limited to "Astrophysical Probes".
- LSST (with complementary instruments) will greatly increase the strength of these probes.
- Mapping the distribution of dark matter tells us where to look with other experiments (cross correlation?).

 Understanding the astrophysical distribution of dark matter is essential for interpreting any particle dark matter detection.





Extra Slides


24

Rate calculation

The differential cross section (for spin-independent interactions) per kilogram of target mass per unit recoil energy is


$$\frac{dR}{dQ} = \frac{\rho_0}{m_\chi} \times \frac{\sigma_0 A^2}{2\mu_p^2} \times F^2(Q) \times \int_{V_m} \frac{f(v)}{v} dv \tag{1}$$

- Dark matter density component, from local and galactic observations
- The unknown particle physics component, hopefully determined by experiment
 - ightharpoonup Proportional to A^2 for most models (written explicitly here)
- ▶ The nuclear part, approximately given by $F^2(Q) \propto e^{-Q/Q_0}$ where $Q_0 \sim \frac{80}{A^{5/3}}$ MeV
- The velocity distribution of dark matter in the galaxy $v_m = \sqrt{Qm_N/2m_r^2}$



Local Dark Matter Velocity





Mao et al. (2012) [1210.2721]